



UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS
General Certificate of Education Advanced Level

MATHEMATICS

9709/71

Paper 7 Probability & Statistics 2 (S2)

October/November 2009

1 hour 15 minutes

Additional Materials: Answer Booklet/Paper
 Graph Paper
 List of Formulae (MF9)



READ THESE INSTRUCTIONS FIRST

If you have been given an Answer Booklet, follow the instructions on the front cover of the Booklet.

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use a soft pencil for any diagrams or graphs.

Do not use staples, paper clips, highlighters, glue or correction fluid.

Answer **all** the questions.

Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.

The use of an electronic calculator is expected, where appropriate.

You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [] at the end of each question or part question.

The total number of marks for this paper is 50.

Questions carrying smaller numbers of marks are printed earlier in the paper, and questions carrying larger numbers of marks later in the paper.

This document consists of **3** printed pages and **1** blank page.



1 2% of biscuits on a production line are broken. Broken biscuits are checked to see whether they are broken. Use a suitable approximation to find the probability that fewer than 4 are broken. [3]

2 The lengths of sewing needles in travel sewing kits are normally distributed with mean μ and standard deviation 1.5 mm. A random sample of 5 is taken. Find the smallest value of μ such that the width of a 95% confidence interval for μ is at most 1 mm. [4]

3 The weights of pebbles on a beach are normally distributed with mean 48.5 grams and standard deviation 12.4 grams.

(i) Find the probability that the mean weight of a random sample of 10 pebbles is greater than 51 grams. [3]

(ii) The probability that the mean weight of a random sample of 10 pebbles is less than 51.6 grams is 0.9332. Find the value of μ . [4]

4 The number of severe floods per year in a certain industry over a 10 year period has followed a Poisson distribution with mean 1.8. Scientists suspect that this has now increased the mean. A hypothesis test, at the 5% significance level, is used to test this suspicion. The number of severe floods, that occur next year will be used for the test.

(i) Show that the rejection region for the test is $X \geq 4$. [5]

(ii) Find the probability of making a Type II error if the true mean number of severe floods is now actually 2.3. [3]

5 The continuous random variable X has probability density function given by

$$f(x) = \begin{cases} k \cos x & 0 \leq x \leq \frac{1}{4}\pi, \\ 0 & \text{otherwise,} \end{cases}$$

where k is a constant.

(i) Show that $k = 4$. [2]

(ii) Find $P(X < 0.4)$. [2]

(iii) Find the upper quartile of X . [3]

(iv) Find the probability that exactly 3 out of 5 random observations are greater than the upper quartile. [2]

6 Photographers often need to take many photographs of a family in order to find a photograph which everyone in the family likes. The number of photographs taken in a trial is taken to be one which everybody likes has mean 15.2. A new photographer claims that in a photograph which everybody likes with fewer photographs taken. To test at the 10% level whether this claim is justified, the numbers of photographs taken by the new photographer with a random sample of 20 families are recorded. The results are summarised below.

(i) Calculate unbiased estimates of the population mean and the variance of the number of photographs taken by the new photographer. [3]

(ii) State null and alternative hypotheses for the test. State the probability that the test results in a Type I error. Say what a Type I error means in the context of the question. [3]

(iii) Carry out the test. [4]

7 The volume of liquid in cans of cola is normally distributed with mean 330 millilitres and standard deviation 5.2 millilitres. The volume of liquid in bottles is normally distributed with mean 500 millilitres and standard deviation 7.1 millilitres.

(i) Find the probability that 3 randomly chosen cans of cola contain less liquid than 2 randomly chosen bottles of tonic water. [5]

(ii) A new drink is made by mixing the contents of 2 cans of cola with 1 bottle of tonic water. Find the probability that the volume of the new drink is more than 900 millilitres. [4]

BLANK PAGE

Permission to reproduce items where third-party ~~copyright~~ protected by copyright is included has been ~~searched where~~ possible. Every reasonable effort has been made by the publisher (UCLES) to ~~right holders~~, but if any items requiring clearance ~~have been~~ included, the publisher will be pleased to make amends at the earliest possible ~~opportuni~~

University of Cambridge International Examinat ~~ions~~ Cambridge Assessment Group. Cambridge Assessment ~~is the~~ brand name of University of Cambridge Local Examinations Syndicate (UCLES), ~~which is a~~ department of the University of Cambridge.