

**MARK SCHEME for the May/June 2011 question paper
for the guidance of teachers**

0606 ADDITIONAL MATHEMATICS

0606/12

Paper 1, maximum raw mark 80

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes must be read in conjunction with the question papers and the report on the examination.

- Cambridge will not enter into discussions or correspondence in connection with these mark schemes.

Cambridge is publishing the mark schemes for the May/June 2011 question papers for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level syllabuses and some Ordinary Level syllabuses.

Page 2	Mark Scheme: Teachers' version	Syllabus
	IGCSE – May/June 2011	0606

Mark Scheme Notes

Marks are of the following three types:

M Method mark, awarded for a valid method applied to the problem. Method marks are not lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate just to indicate an intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem in hand, e.g. by substituting the relevant quantities into the formula. Correct application of a formula without the formula being quoted obviously earns the M mark and in some cases an M mark can be implied from a correct answer.

A Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated method mark is earned (or implied).

B Accuracy mark for a correct result or statement independent of method marks.

- When a part of a question has two or more “method” steps, the M marks are generally independent unless the scheme specifically says otherwise; and similarly when there are several B marks allocated. The notation DM or DB (or dep*) is used to indicate that a particular M or B mark is dependent on an earlier M or B (asterisked) mark in the scheme. When two or more steps are run together by the candidate, the earlier marks are implied and full credit is given.
- The symbol \surd implies that the A or B mark indicated is allowed for work correctly following on from previously incorrect results. Otherwise, A or B marks are given for correct work only. A and B marks are not given for fortuitously “correct” answers or results obtained from incorrect working.
- Note: B2 or A2 means that the candidate can earn 2 or 0.
B2, 1, 0 means that the candidate can earn anything from 0 to 2.

Page 3	Mark Scheme: Teachers' version	Syllabus
	IGCSE – May/June 2011	0606

The following abbreviations may be used in a mark scheme or used on the scripts:

AG	Answer Given on the question paper (so extra checking is needed to ensure that the detailed working leading to the result is valid)
BOD	Benefit of Doubt (allowed when the validity of a solution may not be absolutely clear)
CAO	Correct Answer Only (emphasising that no "follow through" from a previous error is allowed)
ISW	Ignore Subsequent Working
MR	Misread
PA	Premature Approximation (resulting in basically correct work that is insufficiently accurate)
SOS	See Other Solution (the candidate makes a better attempt at the same question)



Penalties

MR –1	A penalty of MR –1 is deducted from A or B marks when the data of a question or part question are genuinely misread and the object and difficulty of the question remain unaltered. In this case all A and B marks then become "follow through $\sqrt{\quad}$ " marks. MR is not applied when the candidate misreads his own figures – this is regarded as an error in accuracy.
OW –1,2	This is deducted from A or B marks when essential working is omitted.
PA –1	This is deducted from A or B marks in the case of premature approximation.
S –1	Occasionally used for persistent slackness – usually discussed at a meeting.
EX –1	Applied to A or B marks when extra solutions are offered to a particular equation. Again, this is usually discussed at the meeting.

Page 4	Mark Scheme: Teachers' version	Syllabus
	IGCSE – May/June 2011	0606

<p>1 $x^2 + (2k + 10)x + (k^2 + 5) = 0$ $(2k + 10)^2 = 4(k^2 + 5)$ $k = -2$</p> <p>(or $\frac{dy}{dx} = 2x + (2k + 10), x = -(k + 5)$ $0 = (k + 5)^2 - (2k + 10)(k + 5) + k^2 + 5$ leading to $k = -2$)</p> <p>(or $(x + A)^2 = x^2 + (2k + 10)x + k^2 + 5$ $A = (k + 5), A^2 = k^2 + 5$ $(k + 5)^2 = k^2 + 5$, leading to $k = -2$)</p> <p>(or by completing the square $y = (x + (k + 5))^2 - (k + 5)^2 + (k^2 + 5)$ $(k + 5)^2 = k^2 + 5$ leading to $k = -2$)</p>	<p>M1 M1 A1 [3] M1 M1 A1 M1 M1 A1 M1 M1 A1</p>	<p>M1 for equating to zero and use $b^2 = 4ac$ M1 for solution</p> <p>M1 for differentiation and attempt to equate to zero. M1 for attempt to substitute in for x in terms of k, for $y = 0$ and for attempt at solution.</p> <p>M1 for approach M1 for equating and attempt at solution</p> <p>M1 for approach M1 for equating last 2 terms to zero and attempt to solve</p>
<p>2 ${}^5C_3 2^2 a^3 = (10)^4 C_2 \frac{a^2}{9}$ $a = \frac{1}{6}$</p>	<p>B1B1 M1 A1 [4]</p>	<p>B1 for ${}^5C_3 2^2 a^3$, B1 for ${}^4C_2 \frac{a^2}{9}$ M1 for a relationship between the 2 coefficients and attempt to solve</p>
<p>3 (a) $k = 2, m = 3, p = 1$ (b) (i) 5 (ii) $\frac{2\pi}{3}$</p>	<p>B3 B1 B1 [5]</p>	<p>B1 for each</p>
<p>There must be evidence of working without a calculator in all parts</p> <p>4 (i) $\frac{(4 + \sqrt{2})(1 - \sqrt{2})}{(1 + \sqrt{2})(1 - \sqrt{2})} = 2\sqrt{2}$ (ii) Area = $\frac{1}{2} \times (4 + 2\sqrt{2}) \times (1 + \sqrt{2})$ $= 4 + 3\sqrt{2}$ (iii) Area = AC^2 $= (4 + 2\sqrt{2})^2 + (1 + \sqrt{2})^2$ $= 27 + 18\sqrt{2}$</p>	<p>M1A1 M1 A1 M1 A1 [6]</p>	<p>M1 for attempt to rationalise and attempt to expand</p> <p>M1 for attempt at area using surd form and attempt to expand</p> <p>M1 for attempt at AC^2 or AC in surd form, with attempt to expand</p>

<p>5 (i) $2\left(\frac{1}{8}\right) - 5\left(\frac{1}{4}\right) + 10\left(\frac{1}{2}\right) - 4$ $= 0$</p> <p>(ii) $(2x-1)(x^2-2x+4)$</p> <p>For (x^2-2x+4), '$b^2 < 4ac$'</p> <p>so only one real root of $x = 0.5$</p>	<p>M1</p> <p>A1</p> <p>M1A1</p> <p>M1</p> <p>A1</p> <p>[6]</p>	<p>M1 for substitution of $x = 0.5$ or at long division</p> <p>M1 attempt to obtain quadratic factor A1 for correct quadratic factor M1 for correct use of discriminant or solution of quadratic equation = 0 A1, all correct with statement of root.</p>
<p>6 (i) $\lg y - 3 = \frac{1}{5}(x-5)$</p> <p>(ii) Either $b = \frac{1}{5}$</p> <p>$y = 10^{\left(\frac{1}{5}x+2\right)}$, $= 10^{\frac{1}{5}x} 10^2$ $a = 100$</p> <p>Or $\lg y = \lg a + \lg 10^{bx}$ $\lg y = \lg a + bx$, $\lg a = 2$ $a = 100$ $b = \frac{1}{5}$</p> <p>Or $10^3 = a(10)^{5b}$ $10^5 = a(10)^{15b}$ $b = \frac{1}{5}$, $a = 100$</p>	<p>B1M1</p> <p>A1</p> <p>B1</p> <p>M1</p> <p>A1</p> <p>M1</p> <p>A1</p> <p>B1</p> <p>M1</p> <p>B1, A1</p> <p>[6]</p>	<p>B1 for gradient, M1 for use of straight line equation</p> <p>B1 for $b = \frac{1}{5}$</p> <p>M1 for use of powers of 10 correctly to obtain a</p> <p>A1 for a</p> <p>M1 for use of logarithms correctly to obtain a</p> <p>A1 for a</p> <p>B1 for $b = \frac{1}{5}$</p> <p>M1 for simultaneous equations involving powers of 10</p> <p>B1 for $b = \frac{1}{5}$, A1 for $a = 100$</p>
<p>7 (i) ${}^{14}C_6 = 3003$</p> <p>(ii) ${}^8C_4 \times {}^6C_2$ $= 1050$</p> <p>(iii) ${}^8C_6 + 6{}^8C_5 = 364$</p>	<p>B1</p> <p>B1B1</p> <p>B1</p> <p>B1B1</p> <p>B1</p> <p>[7]</p>	<p>B1 for 8C_4 or 6C_2</p> <p>B1 for \times by 6C_2 or 8C_4</p> <p>B1 for 1050</p> <p>B1 for 8C_6 or equivalent</p> <p>B1 for $6{}^8C_5$ or equivalent</p> <p>B1 for 364</p>

<p>8 (i) </p> <p>(ii) (1, -9)</p> <p>(iii) </p>	<p>B1 B1 B1 B1</p> <p>B1</p> <p>√B1 B1</p> <p>[7]</p>	<p>B1 for $x = -0.5$ B1 for $x = 2.5$ B1 for $y = -5$ B1 for shape</p> <p>B1</p> <p>√B1 on shape from (i) B1 for a completely correct sketch</p>
<p>9 (i) $\triangle OBA: \theta + 2\left(\frac{\theta}{3}\right) = \pi$</p> <p>(ii) $9\pi = r \times \frac{3\pi}{5}$ $r = 15$</p> <p>(iii) Area = $\left(\frac{1}{2} \times 15^2 \times \frac{3\pi}{5}\right) - \left(\frac{1}{2} \times 15^2 \times \sin \frac{3\pi}{5}\right)$ = 105</p>	<p>M1 A1</p> <p>M1 A1</p> <p>M1M1 A1</p> <p>[7]</p>	<p>M1 for using angles in an isosceles triangle</p> <p>M1 for use of $s = r\theta$</p> <p>M1 for use of $\frac{1}{2}r^2\theta$ or $\frac{1}{2}rs$ M1 for use of $\frac{1}{2}r^2 \sin \theta$ or other correct method</p>
<p>10 (i) $\begin{pmatrix} 29 \\ -13 \end{pmatrix} - \begin{pmatrix} 5 \\ -6 \end{pmatrix} = \begin{pmatrix} 24 \\ -7 \end{pmatrix}$ Magnitude = 25, unit vector $\frac{1}{25} \begin{pmatrix} 24 \\ -7 \end{pmatrix}$</p> <p>(ii) $2\overline{AC} = 3\overline{AB}$ or $2\overline{AB} + 2\overline{BC} = 3\overline{AB}$ leading to $\overline{AC} = \begin{pmatrix} 36 \\ -10.5 \end{pmatrix}$ $\overline{OC} = \overline{OA} + \overline{AC}$ or $\overline{OB} - \overline{OA} = 2\overline{OC} - 2\overline{OB}$ leading to $\overline{OC} = \begin{pmatrix} 41 \\ -16.5 \end{pmatrix}$ (equivalent methods acceptable)</p>	<p>M1</p> <p>M1 A1</p> <p>M1</p> <p>M1</p> <p>A1 A1</p> <p>[7]</p>	<p>M1 for subtraction</p> <p>M1 for attempt to find magnitude of their vector</p> <p>M1 for attempt to find \overline{AC} – may be part of a larger method</p> <p>M1 for attempt to find \overline{OC}</p> <p>A1 for each</p>

<p>11 (i) $2 \cos^2 x - 5 \cos x - 3 = 0$ $(2 \cos x + 1)(\cos x - 3) = 0$ leading to $\sin x = \frac{1}{3}$, $x = 19.5^\circ, 160.5^\circ$</p> <p>(ii) $\tan 2y = \frac{5}{4}$ $2y = 51.34^\circ, 231.34^\circ$ $y = 25.7^\circ, 115.7^\circ$</p> <p>(iii) $\left(z + \frac{\pi}{6}\right) = \frac{2\pi}{3}, \frac{4\pi}{3}$ $z = \frac{2\pi}{3} - \frac{\pi}{6} \quad \left(\frac{4\pi}{3} - \frac{\pi}{6}\right)$ $z = \frac{\pi}{2}, \frac{7\pi}{6}$ allow 1.57, 3.67</p>	<p>M1A1 DM1 A1√A1 M1 M1 A1,√A1 M1 A1, A1 [12]</p>	<p>M1 for use of correct identity or to get in terms of $\sin x$ DM1 for attempt to solve $\sqrt{180^\circ}$ – their x M1 for attempt to get in terms of \tan M1 for dealing correctly with double angle $\sqrt{90^\circ}$ their y M1 for dealing with order correctly and attempt to solve</p>
<p>12 EITHER</p> <p>(i) $\frac{dy}{dx} = 9x^2 + 4x - 5$ when $x = -1$, $\frac{dy}{dx} = 0$ tangent $y = 5$, $A (0, 5)$</p> <p>(ii) $B (0, 1)$ At B, $\frac{dy}{dx} = -5$ normal $y - 1 = \frac{1}{5}x$ $C (-5, 0)$ At D $\frac{1}{5}x + 1 = 5$, $D (20, 5)$ Area = $\frac{1}{2} \times 20 \times 5$, = 50</p>	<p>M1 DM1 A1 B1 M1A1 M1A1 M1 A1 [10]</p>	<p>M1 for differentiation and substitution of $x = -1$ DM1 for attempt at equation of tangent and coordinates of A B1 for B M1 for attempt at normal and C, must be from differentiation and using correct point M1 for attempt to obtain D, equating normal and tangent equations M1 for valid attempt at area</p>

Page 8	Mark Scheme: Teachers' version	Syllabus
	IGCSE – May/June 2011	0606

<p>12 OR</p> $\frac{dy}{dx} = 3x^2 - 12x + 9$ <p>When $\frac{dy}{dx} = 0, x = 1, 3$</p> <p style="text-align: right;">$P(1, 4)$</p> $\text{Area} = 8 - \int_1^3 x^3 - 6x^2 + 9x \, dx$ $= 8 - \left[\frac{x^4}{4} - 2x^3 + \frac{9x^2}{2} \right]_1^3$ $= 8 - \frac{27}{4} + \frac{11}{4}$ $= 4$	<p>M1</p> <p>M1 A1 A1</p> <p>√B1M1</p> <p>A2,1,0</p> <p>DM1</p> <p>A1</p> <p>[10]</p>	<p>M1 for differentiation and equating to zero can be using a product</p> <p>M1 for attempt to solve A1 for both x values A1 for y coordinate</p> <p>√B1 on y coordinate for area of rectangle M1 for attempt to integrate</p> <p>-1 each error</p> <p>DM1 for application of limits</p>
--	---	---