

## UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS International General Certificate of Secondary Education

CANDIDATE NAME					
CENTRE NUMBER		CANDIDATE NUMBER			
CAMBRIDGE INTERNATIONAL MATHEMATICS 0607/22					
CAMBRIDGE	INTERNATIONAL MATHEMATICS		0607/22		
CAMBRIDGE Paper 2 (Exten			0607/22 May/June 2012		
Paper 2 (Exten			May/June 2012		

## READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

Do not use staples, paper clips, highlighters, glue or correction fluid.

You may use a pencil for any diagrams or graphs.

DO NOT WRITE IN ANY BARCODES.

Answer all the questions.

## CALCULATORS MUST NOT BE USED IN THIS PAPER.

All answers should be given in their simplest form.

You must show all the relevant working to gain full marks and you will be given marks for correct methods even if your answer is incorrect.

The number of marks is given in brackets [] at the end of each question or part question.

The total number of marks for this paper is 40.

For Examiner's Use

This document consists of **8** printed pages.




## **Formula List**

For the equation	$ax^2 + bx + c = 0$	$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$
Curved surface area, A, of cylir	nder of radius r, height h.	$A = 2\pi rh$
Curved surface area, <i>A</i> , of cone	e of radius r, sloping edge l.	$A = \pi r l$
Curved surface area, A, of sphe	re of radius <i>r</i> .	$A = 4\pi r^2$
Volume, $V$ , of pyramid, base an	rea A, height h.	$V=\frac{1}{3}Ah$
Volume, $V$ , of cylinder of radiu	s $r$ , height $h$ .	$V = \pi r^2 h$
Volume, $V$ , of cone of radius $r$ ,	height <i>h</i> .	$V = \frac{1}{3}\pi r^2 h$
Volume, $V$ , of sphere of radius	r.	$V = \frac{4}{3}\pi r^3$
	C	$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$ $a^2 = b^2 + c^2 - 2bc \cos A$ $\operatorname{Area} = \frac{1}{2}bc \sin A$

For Answer **all** the questions. Examiner's Use(a) Find the value of  $49^{-\frac{1}{2}}$ . 1 Answer(a) [1] **(b)** When  $x^{-2} = 4$  write down the values of *x*. Answer(b) x = or x =[2] 2 (a) Factorise  $6x^2 - x - 2$ . Answer(a) [2] (b) Solve the equation  $6x^2 - x - 2 = 0$ . Answer(b) x = or x =[1]

$\mathbf{p} = \begin{pmatrix} 2\\ 3 \end{pmatrix} \qquad \mathbf{q} = \begin{pmatrix} 1\\ 1 \end{pmatrix}$	$\begin{pmatrix} -3 \\ 5 \end{pmatrix}$				F Exam U
Find					
(a) $2p - 3q$ ,					
					[0]
				Answer(a)	[2]
(b) <b> </b> p <b> </b> .					
				Answer(b)	 [2]
<b>F</b> ' 1.1	o terms i	n this sequ	ence.		
Find the next two					
	2,	6,	15,	31,	
		6,	15,	31,	
		6,	15,	31,	
		6,	15,	31,	
		6,	15,	31,	
		6,	15,	31,	
		6,	15,		 [2]
		6,	15,		 [2]
		6,	15,		 [2]

4



© UCLES 2012

7 .1

7 Solve the simultaneous equations.  

$$3p + 4q = 7$$

$$5p + 6q = 10$$
  
Answer  $p = \dots$ 

$$q = \dots$$


$$q = \dots$$

$$[4]$$
8 y varies directly as  $x^2$ , where x is a positive integer.

When x = 3, y = 108.

Calculate the value of *x* when y = 300.

[3] Answer x =.....



During one training session he

- swims 1 km in 15 minutes,
- cycles 20 km at a speed of 20 km/h,
- runs at a speed of 8 km/h for 45 minutes.

Calculate Joe's average speed for the training session. Give your answer in kilometres per hour.

Answer km/h [3]

10 Solve the equation.

$$\frac{x+3}{7} - \frac{3(x-1)}{14} = 1$$

Answer x = [3]

Questions 11 and 12 are on the next page.

For Examiner's Use 8

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.